


BLOOD SAMPLE COLLECTION KIT INSTRUCTIONS

Please read these instructions carefully before collecting your sample.



VICTUS<sup>88</sup> KIT CONTENTS




Lancet x4



Gauze x2



Wipe x2




Bandage x4



4-sampler  
Clamshell x1



Silver Specimen  
Bag x1




FedEx Pack  
w/Label x1



Label for  
Clamshell



FIRST, UNPACK AND PREPARE YOUR KIT CONTENTS




Twist off  
lancet caps



Tear open silver  
specimen bag




Remove  
Clamshell



Open  
Clamshell



Unwrap gauze  
& wipes




Unpeel back of  
bandages


STEP-BY-STEP INSTRUCTIONS




1 **Wash your hands**  
with warm, soapy water.




2 **Warm your hands**  
by rubbing them together. You  
may also warm your hands by  
walking, jumping, or any other  
method you prefer. **Warming  
your hands is IMPORTANT for  
good blood flow.**




3 **Select the finger you will use**  
to provide the blood sample.  
See illustration for recommend-  
ed locations off-center of finger  
midline. Use alcohol wipes, if  
desired, to clean sample area.




4 **Prick your finger**  
by laying hand on a hard sur-  
face. Position the lancet on the  
chosen location. Push down on  
the end of lancet until a click  
is heard.



5 **Wipe away first blood drop**  
with the gauze. A blood drop may  
take up to 20 seconds to form. If  
needed, gently milk pricked  
finger upward until drop forms.



**note** - you may find that wiping away another drop of  
blood after filling a few sampling tips helps with blood  
flow.



You may apply sampling tips  
to the blood drop multiple  
times to fill.  
Always point sampling tip  
towards floor when touching  
it to blood drop on your  
finger as shown above. Do  
NOT drip blood onto tip.

6 **Collect sample by touching**  
one of the four sampling tips to the  
blood drop. First, watch it turn fully  
red. Next, count 2 seconds then  
slowly remove the tip from your  
finger. Repeat this process with the  
other three sampling tips.

**ensure sampling tips are filled correctly**  
Under-sampling occurs when the sampling tip is  
removed from the finger too soon. No white should remain.

If so, repeat steps 4-6 with same tip until it turns fully red.



over-sampled




under-sampled



correctly sampled


Over-sampling occurs when blood is dripped onto the  
sampling tip from above. Always touch sampling tip to  
blood drop as shown in step 6.




7 **Set Clamshell device down**  
facing up on hard surface and  
apply bandage to your finger.



8 **Close & label Clamshell device**  
by lifting the sides to meet at the  
top. Press together until a click is  
heard. Place one label from bottom  
of requisition form on the clamshell.  
Write collection date on label.



9 **Insert Clamshell device**  
into the specimen bag and seal shut.  
**Ensure the desiccant is still in the  
bag.**



10 **Insert specimen bag**  
into FedEx shipping envelope along  
with requisition form. Attach FedEx  
shipping label and drop off at nearest  
FedEx store location.

Intended Use

A single use, non sterile device used as a container to collect, store and transport blood specimen for research use.



Phone: 618.690.9555  
Fax: 618.400.1220



1616 Eastport Plaza Drive  
Collinsville, IL 62234



info@biovisiondx.com  
www.victus88.com